Please let me know if any of the problems are unclear or have typos.

**Exercise 1.1.** Suppose that X is a topological space. Define Homeo(X) to be the set of homeomorphisms  $f: X \to X$ . Show that Homeo(X) is a group if we take the group operation to be function composition.

Give an example of a space X where Homeo(X) is the trivial group.

**Exercise 1.2.** Show that the relation  $X \cong Y$  of being homeomorphic is an equivalence relation on topological spaces. Now consider the capital letters of the alphabet A, B, C, ... in a sans serif font. Each of these gives a graph in the plane. Sort these into homeomorphism classes. (The partition may depend on the font! In particular, K can be tricky.)

**Exercise 1.3.** We equip  $[0,1) \subset \mathbb{R}$  and  $S^1 \subset \mathbb{C}$  with their usual subspace topologies. Consider the map  $p: [0,1) \to S^1$  given by  $p(t) = \exp(2\pi i t)$ . Show that p is a continuous bijection. Show that p is not a homeomorphism.

**Exercise 1.4.** We equip  $[0,1] \subset \mathbb{R}$  and  $S^1 \subset \mathbb{C}$  with their usual subspace topologies. Show that the quotient space

$$X = [0, 1] / 0 \sim 1$$

is homeomorphic to  $S^1$ .

**Exercise 1.5.** For three of the following pairs (X, Y) show that X is not homeomorphic to Y.

- The graph X and the graph Y.
- (0,1) and [0,1]: the open and closed intervals.
- $S^1$  and [0,1]: the circle and the closed interval.
- $S^1$  and  $S^2$ : the circle and the sphere.
- $\mathbb{R}^1$  and  $\mathbb{R}^2$ : the line and the plane.
- $\mathbb{R}^2$  and  $\mathbb{R}^3$ : the plane and three-space (harder).
- $S^2$  and  $T^2 = S^1 \times S^1$ : the sphere and the torus (harder).

**Exercise 1.6.** Suppose X and Y are topological spaces. We call a function  $f: X \to Y$  an *embedding* if f is a homeomorphism from X to f(X), equipped with the subspace topology. Give an example of a space X that does not embed in  $\mathbb{R}^n$ , for any n.