The required problems are Exercises 6.1, 6.3, and 6.4. Please let me know if any of the problems are unclear or have typos.

**Exercise 6.1.** Recall that  $T^2 = S^1 \times S^1$  is the two-torus; informally  $T^2$  is the surface of a donut. Fix any point  $x \in T^2$ ; show that  $T^2 - \{x\}$  deformation retracts to the figure-eight graph. Illustrate your proof with useful figures.

**Exercise 6.2.** [A version of Exercise 14, page 39, of Hatcher's book.] Given topological spaces X and Y we equip  $Z = X \times Y$  with the product topology. Let  $p: X \times Y \to X$  be projection to the first factor; that is p(a, b) = a. Define  $q: X \times Y \to Y$  to be projection to the second factor.

Fix  $x \in X$  and  $y \in Y$ . Prove that the homomorphism

$$p_* \times q_* \colon \pi_1(X \times Y, (x, y)) \to \pi_1(X, x) \times \pi_1(Y, y)$$

is an isomorphism. (Essentially you are being asked to carefully reprove Proposition 1.12, using the notion of projections.)

**Exercise 6.3.** The *real projective space*  $\mathbb{RP}^n$  is the space of lines through the origin in  $\mathbb{R}^{n+1}$ . We equip  $\mathbb{RP}^n$  with its usual topology, coming from the round metric; the distance between distinct lines  $L, M \subset \mathbb{R}^{n+1}$  is the smaller of the two angles made by L and M in the plane they span.

- Exhibit a two-fold covering map  $p: S^n \to \mathbb{RP}^n$ .
- Deduce that  $\pi_1(\mathbb{RP}^n) \cong \mathbb{Z}/2\mathbb{Z}$ , when  $n \ge 2$ .
- Briefly discuss the cases of n = 0 and n = 1. Give pictures.

**Exercise 6.4.** Suppose that  $p: \widetilde{X} \to X$  is a *d*-fold covering map and that  $\widetilde{X}$  is pathconnected. Prove that Deck(p) has at most *d* elements. Give examples which do and which do not realize this bound.

**Exercise 6.5.** [Hard.] Let X and Y be copies of the two-sphere and choose distinct points  $p, p' \in X$  and  $q, q' \in Y$ . Define

$$Z = X \sqcup Y / p \sim q, \ p' \sim q'$$

to be the quotient space. That is, Z is obtained from the disjoint union of X and Y by identifying p with q and p' with q'. Draw a picture of Z. Compute  $\pi_1(Z)$  and carefully justify your reasoning.